

CP-VIOLATION^a

J. Bernabéu

*Departamento de Física Teórica, Univ. de Valencia
46100 Burjassot, Valencia, Spain*

These lectures cover different aspects of the subject of CP-Violation, from its description inside the Standard Model to the phenomenological analysis of the $K^0 - \bar{K}^0$ system and the prospects for its manifestation in B-physics.

Contents

1. CP in the Standard Model
 - 1.1 Discrete symmetries: P,C
 - 1.2 Framework of local field theories
 - 1.3 CPT Invariance: antiparticles
 - 1.4 The Standard Electroweak Model
 - 1.5 Yukawa Couplings
 - 1.6 Spontaneous Symmetry Breaking
 - 1.7 The Quark Mixing Matrix
 - 1.8 CP-Violation with 3 families
2. The $K^0 - \bar{K}^0$ System
 - 2.1 Discovery of CP-Violation
 - 2.2 Meson-Antimeson Mixing
 - 2.3 Indirect CP-Violation
 - 2.4 Isospin Decomposition
 - 2.5 Experiments for $K_L \rightarrow 2\pi$
 - 2.6 CKM Quark Mixing Matrix
 - 2.7 Coherent Decay of $(K^0, \bar{K}^0)$
 - 2.8 Time Integrated Rates
3. CP-Violation and B-Physics
 - 3.1 Physics Motivation
 - 3.2 Principle of the CP-violation measurement
 - 3.3 Rate Asymmetries
 - 3.4 Outlook

^aLectures delivered at the XXIV International Meeting on Fundamental Physics, Playa de Gandía (1996)

1 CP in the Standard Model

1.1 Discrete symmetries: P , C

Conservation Laws in Physics are due to invariance of forces under symmetry transformations.

In particular, there are conservation laws corresponding to discrete transformations:

(i) Reflection in space or Parity (P). In three dimensions, it means that the mirror image of an experiment yields the same results as the original experiment. This implies

P -invariance means that “left” and “right” cannot be defined in an absolute sense.

(ii) Particle-antiparticle conjugation (C), i.e., C transforms each particle into its antiparticle, without touching its space-time properties. Similarly,

C -invariance of laws means that experiments in a world of antimatter will give identical results to the ones in our world.

As an example, C -invariance for electromagnetic interactions implies that the atom of anti-hydrogen, recently discovered¹ and manipulated at CERN-LEAR, should show the same spectral lines as those of the hydrogen atom.

Until the work of Lee and Yang² in 1956 it was assumed that all elementary processes are invariant under the separate symmetries P and C . Subsequent experiments in nuclear β -decay, $\pi^\pm$ decays and $\mu^\pm$ decays demonstrated the violation of P and C invariance under weak interactions. The application of these symmetries to $\pi^\pm$ decays is shown in Fig.1, where the configurations for momenta and helicities of the decay products are exhibited for the $\pi^- \rightarrow \mu^- \bar{\nu}_\mu$ process and its P , C and CP transformed processes. Nature selects that only the μ^- -helicity $+1$ in π^- -decay and the μ^+ -helicity -1 in π^+ -decay are present, so P and C symmetries are violated whereas CP is a good symmetry.




Figure 1: C,P and CP applied to $\pi \rightarrow \mu\nu$

We conclude that

The combined operation CP connects two physical (existing in Nature!) processes.

CP-invariance was considered³ to be replacing the separate P and C invariances of weak interaction. Not only for charge current weak interactions: neutral current weak interactions, mediated by the Z vector boson, are known at present to violate P and C too.

P -Violation by neutral currents was beautifully demonstrated by the left-right asymmetry in deep inelastic scattering of incoming electrons on a deuterium target at SLAC⁴ and by parity violation in atoms⁵, where the interference of the electromagnetic interaction with the P -odd neutral current interaction operates. More recently, at the peak of the Z , the left-right asymmetry for electron-positron collisions at SLC⁶ measures the P -odd combination of neutral current couplings of the electron, whereas the P -odd τ polarization⁷ has been measured by LEP1. The forward-backward asymmetry in the process $e^+e^- \rightarrow f\bar{f}$ has been observed⁷ first by LEP1 for both leptons and quarks in the final state and it demonstrates C -violation in the neutral current interaction.

1.2 Framework of local field theories

In the framework of local field theories, one introduces a lagrangian density

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{int}$$

where $\mathcal{L}_0$ is the free part, describing noninteracting particles, and $\mathcal{L}_{int}$ gives how these particles interact. For N fields $\phi_j(x)$, we assume

$$\mathcal{L}(x) = \mathcal{L}(\phi_j(x), \partial^\mu \phi_j(x)); \quad j = 1, 2, \dots, N$$

For the discussion of symmetries, the fundamental quantity is the ACTION

$$A = \int d^4x \mathcal{L}(x) = A_0 + A_{int}$$

If the action A is (is not) invariant under a symmetry operation, then the symmetry is a good (broken) one.

We define the symmetry operation P, C for the free fields entering $\mathcal{L}_0$ and $\mathcal{L}_{int}$: A field and its P-or C-image satisfy the SAME EQUATION OF MOTION. One should keep in mind that the fields in the Lagrangian density may or may not correspond to physical fields (those describing particles with well defined mass, lifetime, ...), a comment which is particularly relevant in theories with spontaneous symmetry breaking.

Parity P ($\vec{x} \rightarrow -\vec{x}$) : $\vec{p} \rightarrow -\vec{p}, \vec{J} \rightarrow \vec{J}$

The free fields transform according to

	P
Scalar field	$\phi(t, \vec{x}) \rightarrow \phi(t, -\vec{x})$
Pseudoscalar field	$P(t, \vec{x}) \rightarrow -P(t, -\vec{x})$
Dirac spinor	$\psi(t, \vec{x}) \rightarrow \gamma_0 \psi(t, -\vec{x})$ $\bar{\psi}(t, \vec{x}) \rightarrow \bar{\psi}(t, -\vec{x}) \gamma_0$
Vector field	$V_\mu(t, \vec{x}) \rightarrow V_\mu(t, -\vec{x})$
Axial Field	$A_\mu(t, \vec{x}) \rightarrow -A^\mu(t, -\vec{x})$

Note that $\mu = 0, 1, 2, 3$ and, for any four vector Q^μ , one has $Q^0 = Q_0$, whereas $Q^k = -Q_k$ for $k = 1, 2, 3$.

The parity of a free field is not observable. It is the interaction which fixes the “relative” parties of the fields, if P is a good symmetry.

For the Dirac spinors, the transformation properties of the bilinears under P are given by

	P
	$(t, \vec{x}) \rightarrow (t, -\vec{x})$
Scalar	$\bar{\psi}_1 \psi_2 \rightarrow \bar{\psi}_1 \psi_2$
Pseudoscalar	$\bar{\psi}_1 \gamma_5 \psi_2 \rightarrow -\bar{\psi}_1 \gamma_5 \psi_2$
Vector	$\bar{\psi}_1 \gamma_\mu \psi_2 \rightarrow \bar{\psi}_1 \gamma_\mu \psi_2$
Axial	$\bar{\psi}_1 \gamma_\mu \gamma_5 \psi_2 \rightarrow -\bar{\psi}_1 \gamma^\mu \gamma_5 \psi_2$
Tensor	$\bar{\psi}_1 \sigma_{\mu\nu} \psi_2 \rightarrow \bar{\psi}_1 \sigma^{\mu\nu} \psi_2$
Pseudotensor	$\bar{\psi}_1 \sigma_{\mu\nu} \gamma_5 \psi_2 \rightarrow \bar{\psi}_1 \sigma^{\mu\nu} \psi_2$

These bilinears are very fundamental objects and appear frequently in physics. From the transformation properties of the fields, written in terms of the annihilation and the creation operators for particles $a(\vec{p}, \lambda)$ and antiparticles $b(\vec{p}, \lambda)$, one can find how the states $|\vec{p}, \lambda\rangle$ transform.

C-conjugation: $a(\vec{p}, \lambda) \rightleftharpoons b(\vec{p}, \lambda)$

The free fields have a Fourier decomposition in terms of the annihilation operator $a(\vec{p}, \lambda)$ for particles plus the creation operator $b^\dagger(\vec{p}, \lambda)$ for antiparticles. Under C-conjugation, the role of the operators $a(\vec{p}, \lambda)$ and $b(\vec{p}, \lambda)$ is interchanged. This exchange does not necessarily mean that the physical states of particles and antiparticles are connected by the C-operation, because the C-symmetry is violated in Nature. As an example, neutrinos and antineutrinos of the same helicity do not exist in Nature.

Under C , the free fields transform as follows

	C
Scalar field	$\phi(x) \rightarrow \phi^+(x)$
Dirac spinor	$\psi(x) \rightarrow C\bar{\psi}^T(x)$ $\bar{\psi}(x) \rightarrow -\psi^T(x)C^{-1}$
Vector field	$V_\mu(x) \rightarrow -V_\mu^+(x)$
Axial field	$A_\mu(x) \rightarrow A_\mu^+(x)$

where C is a 4×4 unitary matrix which satisfies the condition

$$C^{-1}\gamma_\mu C = -\gamma_\mu^T$$

as a result of imposing that the C-conjugated Dirac field satisfies the same free field equation of motion as the field itself. The matrix C can be realized by the choice $C = i\gamma^2\gamma^0$.

For a Real Vector Field, such as the photon γ or the neutral rho ρ^0 , charged conjugation C is well defined with eigenvalue $C_\gamma = -1$.

The spinor bilinears are transformed under C as.

	$x \rightarrow x$
Scalar	$\bar{\psi}_1 \psi_2 \rightarrow \bar{\psi}_2 \psi_1$
Pseudoscalar	$\bar{\psi}_1 \gamma_5 \psi_2 \rightarrow \bar{\psi}_2 \gamma_5 \psi_1$
Vector	$\bar{\psi}_1 \gamma_\mu \psi_2 \rightarrow -\bar{\psi}_2 \gamma_\mu \psi_1$
Axial	$\bar{\psi}_1 \gamma_\mu \gamma_5 \psi_2 \rightarrow \bar{\psi}_2 \gamma_\mu \gamma_5 \psi_1$
Tensor	$\bar{\psi}_1 \sigma_{\mu\nu} \psi_2 \rightarrow -\bar{\psi}_2 \sigma_{\mu\nu} \psi_1$
Pseudotensor	$\bar{\psi}_1 \sigma_{\mu\nu} \gamma_5 \psi_2 \rightarrow -\bar{\psi}_2 \sigma_{\mu\nu} \gamma_5 \psi_1$

CP-operation: $(t, \vec{x}) \rightarrow (t, -\vec{x})$

Scalar	$\bar{\psi}_1 \psi_2 \rightarrow \bar{\psi}_2 \psi_1$
Pseudoscalar	$\bar{\psi}_1 \gamma_5 \psi_2 \rightarrow -\bar{\psi}_2 \gamma_5 \psi_1$
Vector	$\bar{\psi}_1 \gamma_\mu \psi_2 \rightarrow -\bar{\psi}_2 \gamma^\mu \psi_1$
Axial	$\bar{\psi}_1 \gamma_\mu \gamma_5 \psi_2 \rightarrow -\bar{\psi}_2 \gamma^\mu \gamma_5 \psi_1$
Tensor	$\bar{\psi}_1 \sigma_{\mu\nu} \psi_2 \rightarrow -\bar{\psi}_2 \sigma^{\mu\nu} \psi_1$
Pseudotensor	$\bar{\psi}_1 \sigma_{\mu\nu} \gamma_5 \psi_2 \rightarrow \bar{\psi}_2 \sigma^{\mu\nu} \gamma_5 \psi_1$

As an example, for an interaction Lagrangian of pseudoscalar-fermion-fermion fields (think in $K_L \rightarrow \mu^+ \mu^-$) given by

$$\mathcal{L}_{int}(x) = \bar{\psi}(x)[a + ib\gamma_5]\psi(x)\phi(x)$$

the process violates parity P , conserves charge conjugation C and violates the combined CP .

1.3 CPT Invariance: antiparticles

A very important property of LOCAL FIELD THEORIES which respect Lorentz invariance is that they are automatically invariant⁸ under the combined operation CPT. This implies that the problem of the invariance under CP is equivalent to that of the invariance under time-reversal T .

Taking into account the role of T to transform the fourvectors x^μ and p^μ and the angular momentum $\vec{J}$.

$$(t, \vec{x}) \xrightarrow{T} (-t, \vec{x}); (p^0, \vec{p}) \xrightarrow{T} (p^0, -\vec{p}); \vec{J} \xrightarrow{T} -\vec{J}$$

the spinor bilinears are transformed under CPT as follows

		CPT	
	$(t, \vec{x})$	$\rightarrow$	$(-t, -\vec{x})$
	$c - \text{number}$	$\rightarrow$	$(c - \text{number})^*$
Scalar	$\bar{\psi}_1 \psi_2$	$\rightarrow$	$\bar{\psi}_2 \psi_1$
Pseudoscalar	$i\bar{\psi}_1 \gamma_5 \psi_2$	$\rightarrow$	$i\bar{\psi}_2 \gamma_5 \psi_1$
Vector	$\bar{\psi}_1 \gamma_\mu \psi_2$	$\rightarrow$	$-\bar{\psi}_2 \gamma_\mu \psi_1$
Axial	$\bar{\psi}_1 \gamma_\mu \gamma_5 \psi_2$	$\rightarrow$	$-\bar{\psi}_2 \gamma_\mu \gamma_5 \psi_1$
Tensor	$\bar{\psi}_1 \sigma_{\mu\nu} \psi_2$	$\rightarrow$	$\bar{\psi}_2 \sigma_{\mu\nu} \psi_1$
Pseudotensor	$i\bar{\psi}_1 \sigma_{\mu\nu} \gamma_5 \psi_2$	$\rightarrow$	$i\bar{\psi}_2 \sigma_{\mu\nu} \gamma_5 \psi_1$

Since 1964⁹ we know that, not only C but, CP is also a broken symmetry. This implies that the definition of the physical states for antiparticles needs the use of the CPT-symmetry.

1.4 The Standard Electroweak Model

The present framework to discuss CP-violation is the Standard Model. The Lagrangian density of the Electroweak Model is of the form

$$\mathcal{L} = \mathcal{L}(f, G) + \mathcal{L}(f, H) + \mathcal{L}(G, H) + \mathcal{L}(G) - V(H)$$

where f = fermions (quarks, leptons), G = gauge bosons ($\vec{W}$ and B), H = the scalar doublet.

The Lagrangian is constructed so that it is invariant under the local (space-time dependent) symmetry group $SU(2) \times U(1)$. Under $SU(2)$, the quark fields transform as doublets, if they are left-handed, and as singlets if they are right-handed. One introduces the multiplets

$$\left\{ \begin{array}{c} q_j \\ q'_j \end{array} \right\}_L \quad ; \quad q_{jR} \ , \ q'_{jR} \quad ; \quad j = 1, 2, \dots, N$$

where the index j is the family index and N denotes the number of families.

The hadronic part of the Lagrangian between fermions and gauge bosons is

$$\begin{aligned} \mathcal{L}(f, G) = \sum_{j=1}^N & \left\{ (\overline{q}, \overline{q'})_{jL} i\gamma^\mu [\partial_\mu - ig_2 \frac{\vec{\sigma}}{2} \cdot \vec{W}_\mu - ig_1 (\frac{1}{6}) B_\mu] \begin{Bmatrix} q \\ q' \end{Bmatrix}_{jL} \right. \\ & \left. + \overline{q}_{jR} i\gamma^\mu [\partial_\mu - ig_1 (\frac{2}{3}) B_\mu] q_{jR} + \overline{q'}_{jR} i\gamma^\mu [\partial_\mu - ig_1 (-\frac{1}{3}) B_\mu] q'_{jR} \right\} \end{aligned}$$

where the numbers in the parenthesis are the eigenvalues of the weak hypercharge Y , the generator of the $U(1)$ group. They are chosen such that $Q \equiv I_3 + Y$ has the eigenvalues $\frac{2}{3}$ for the up-type quarks and $-\frac{1}{3}$ for the down-type quarks.

The Lagrangian density $\mathcal{L}(f, G)$ violates both P and C symmetries. For P , we observe that the interaction of left-handed quarks is different from the interaction of right-handed quarks. This P -non invariance remains even after the lagrangian is rewritten in terms of the physical fields. The simultaneous presence of vector and axial interactions leads to the violation of C -symmetry, because the vector density is odd, whereas the axial density is even. However, $\mathcal{L}(f, G)$ is CP -invariant¹⁰.

1.5 Yukawa Couplings

The Lagrangian density $\mathcal{L}(G, H)$ between the gauge bosons and the scalars is both P - and C -symmetric.

All CP -violation, in the Electroweak Standard Model, originates from $\mathcal{L}(f, H)$.

The hadronic part of the Yukawa interaction between fermions and scalars is given by

$$\begin{aligned} \mathcal{L}(f, H) = & \sum_{j,k=1}^N \left\{ Y_{jk} (\overline{q}, \overline{q'})_{jL} \begin{Bmatrix} \phi^{(0)*} \\ -\phi^{(-)} \end{Bmatrix} q_{kR} \right. \\ & \left. + Y'_{jk} (\overline{q}, \overline{q'})_{jL} \begin{Bmatrix} \phi^{(+)} \\ \phi^{(0)} \end{Bmatrix} q'_{kR} + \text{h.c.} \right\} \end{aligned}$$

The Yukawa couplings Y_{jk}, Y'_{jk} are arbitrary complex numbers, because $\mathcal{L}(f, H)$ is manifestly invariant under the gauge group $SU(2) \times U(1)$ symmetry. The Lagrangian density $\mathcal{L}(f, H)$ involves scalar and pseudoscalar interactions, so it violates P, C and CP symmetries. However, in this case it is of relevance to discuss the symmetry properties after spontaneous symmetry breakdown.

1.6 Spontaneous Symmetry Breaking

Under the spontaneous symmetry breaking, in the unitary gauge, the complex field $\phi^{(0)}$ is shifted and becomes real and the field $\phi^{(+)}$ vanishes:

$$\left. \begin{aligned} \phi^{(0)} &\rightarrow \frac{1}{\sqrt{2}}(v + H) \\ \phi^{(+)} &\rightarrow 0 \end{aligned} \right\} \begin{aligned} v &\equiv \text{v.e.v.} \\ H &\equiv \text{Higgs field} \end{aligned}$$

From the four degrees of freedom of the scalar doublet, three fields are "eaten" by the longitudinal components of the $W^\pm$ and Z bosons, which become massive.

After spontaneous symmetry breaking, what remains for $\mathcal{L}(f, H)$ is

$$\mathcal{L}(f, H) \xrightarrow{SSB} - \sum_{j,k=1}^N \{ m_{jk} \bar{q}_{jL} q_{kR} + m'_{jk} \bar{q}'_{jL} q'_{kR} + \text{h.c.} \} \left[1 + \frac{1}{v} H \right]$$

where the quantities

$$m_{jk} = -\frac{v}{\sqrt{2}} Y_{jk} \quad , \quad m'_{jk} = -\frac{v}{\sqrt{2}} Y'_{jk}$$

are the complex QUARK MASS MATRICES. Their dimension is $N \times N$, N being the number of families. This Lagrangian violates the discrete symmetries P , C and CP :

P	is violated if	$[m], [m']$	are not hermitian.
C	is violated if	$[m], [m']$	are not symmetric.
CP	is violated if	$[m], [m']$	are not real.

However, we should keep in mind that we are, so far, dealing with non-physical fields. In order to find the physical fields we must diagonalize the quark mass matrices $[m], [m']$. We know that any square matrix, hermitian or not, can be diagonalised by means of two unitary matrices:

$$\begin{aligned} U_L \quad m U_R^+ &= D \quad \equiv \text{Diag}(m_u, m_c, m_t) \\ U_L' \quad m' U_R'^+ &= D' \quad \equiv \text{Diag}(m_d, m_s, m_b) \end{aligned}$$

We can discover the meaning of U_L as

$$U_L m U_R^+ U_R m^+ U_L^+ = U_L m m^+ U_L^+ = D^2$$

the unitary matrix which diagonalizes the hermitian matrix mm^+ . Similarly, U_R is the unitary matrix diagonalizing m^+m . Analogous relations can be derived for the down-quark sector, putting primes on the U 's.

In order to identify the physical fields, we write from the Lagrangian density

$$\begin{aligned}\bar{q}_{jL}m_{jk}q_{kR} &\equiv \bar{q}_L m q_R = \bar{q}_L U_L^+ U_L m U_R^+ U_R q_R \\ &= \overline{U_L q_L} D U_R q_R = \overline{U_L q_L} \begin{pmatrix} m_u & 0 & 0 \\ 0 & m_c & 0 \\ 0 & 0 & m_t \end{pmatrix} U_R q_R\end{aligned}$$

Thus the physical fields are

$$q_L^{phys} = U_L q_L = U_L \begin{Bmatrix} u_L \\ c_L \\ t_L \end{Bmatrix} \quad ; \quad q_L'^{phys} = U_L' q_L' = U_L' \begin{Bmatrix} d_L \\ s_L \\ b_L \end{Bmatrix}$$

with similar relations valid for the R-handed quarks. The Yukawa Lagrangian becomes

$$\begin{aligned}\mathcal{L}^{phys}(f, H) &= -(1 + \frac{H}{v})[m_u \bar{u} \frac{1+\gamma_5}{2} u + m_c \bar{c} \frac{1+\gamma_5}{2} c + ... \\ &\quad + m_d \bar{d} \frac{1+\gamma_5}{2} d + m_s \bar{s} \frac{1+\gamma_5}{2} s + ...] + \text{h.c.} \\ &= -(1 + \frac{H}{v})[m_u \bar{u} u + m_c \bar{c} c + ... \\ &\quad + m_d \bar{d} d + m_s \bar{s} s + ...]\end{aligned}$$

The essential feature of $\mathcal{L}^{phys}(f, H)$ is that it conserves, separately, P and C , and thus also CP and T . The lesson to be learnt¹¹ from this exercise is that the "apparent" symmetry properties of a Lagrangian density need not have anything to do with Physics, when this Lagrangian is expressed in terms of unphysical fields.

At this point, it is of interest to summarize our findings for the Standard Model:

- 1) Given the quark mass matrices m, m' , we need four unitary matrices $U_L, U_R ; U'_L, U'_R$ to diagonalize them.
- 2) The unitary diagonalizing matrices allow to find the relations between physical (eigenfields of mass) and nonphysical (fields with definite transformation properties under the gauge group) fields.
- 3) The Lagrangian density $\mathcal{L}^{phys}(f, H)$, in terms of physical fields, has no traces of the unitary U, U' matrices.
- 4) The physical Higgs-fermion interaction conserves P, C and it is Flavour Diagonal.
- 5) The above properties 1) $\rightarrow$ 4) could change if one introduces Physics Beyond the Standard Model.

1.7 The Quark Mixing Matrix

Now that we have the term $\mathcal{L}(f, H)$, in terms of the physical quark fields, we turn our attention to the other term of the Lagrangian density which needs to be expressed in terms of the physical quark fields, i.e. $\mathcal{L}(f, G)$. Consider first any "neutral current" term, i.e., any term which involves only either the up-kind quarks or the down-kind quarks but not both. As an example.

$$\begin{aligned}
\bar{q}_{jR} \gamma^\mu [\partial_\mu - ig_1(\frac{2}{3})B_\mu] q_{jR} &= \\
&= \bar{q}_R^{phys} U_R \gamma^\mu [\partial_\mu - ig_1(\frac{2}{3})B_\mu] U_R^\dagger q_R^{phys} \\
&= \bar{q}_R^{phys} \gamma^\mu [\partial_\mu - ig_1(\frac{2}{3})B_\mu] q_R^{phys}
\end{aligned}$$

As a consequence of unitarity $U_R U_R^\dagger = 1$ and U_R again "disappears": neutral Current terms are Flavour Conserving. In the Standard Model there are some "miracles" which happen "naturally" [see¹²]:

- i) Neutral Currents all conserve CP.
- ii) There are No Flavour-Changing-Neutral-Currents, due to two facts
 - these terms are helicity conserving
 - the four matrices U, U' (with indices L or R) are unitary.

This absence of flavour changing neutral currents is usually referred to as the GIM mechanism¹³. As there are stringent limits, from experiment, on FCNC, the absence of such currents in the Standard Model is one of its great successes. One can note that Physics Beyond the Standard Model generally encounters difficulties in this respect.

What remains to be treated in $\mathcal{L}(f, G)$ is the charged current term. Such terms are helicity conserving, but do mix the up-type and the down-type quarks, simply because $W^\pm$ carry one unit of charge. Since the charged currents only involve left-handed quarks the matrices U_R, U'_R do not enter, but $U_L U_L'^+ \neq I$.

The charged current terms are given by

$$\begin{aligned} [W_\mu^1 - iW_\mu^2] \bar{q}_L \gamma^\mu q'_L + \text{h.c.} &= [W_\mu^1 - iW_\mu^2] \bar{q}_L^{phys} \gamma^\mu U_L U_L'^+ q_L'^{phys} + \text{h.c.} = \\ [W_\mu^1 - iW_\mu^2] \bar{q}_L^{phys} \gamma^\mu V q_L'^{phys} + \text{h.c.} &\equiv [W_\mu^1 - iW_\mu^2] J_{cc}^\mu + \text{h.c.} \end{aligned}$$

where V is a unitary matrix and

$$J_{cc}^\mu \equiv [\bar{u}, \bar{c}, \bar{t}]_L \gamma^\mu \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{Bmatrix} d \\ s \\ b \end{Bmatrix}_L$$

The matrix V is the Quark Mixing Matrix.

We note the following properties:

- 1) All CP-Violation is in V , together with Flavour Mixing.
- 2) Charged Currents violate P, C maximally.
- 3) CP-symmetry would require V real, modulus unmeasurable phases.

1.8 CP-Violation with 3 Families

What matters in field theory is not the absolute phases but the relative phases of different fields. Which phases in V are measurable ? Under rephasing of the L-quark fields, one finds

$$V_{\alpha j} \rightarrow e^{i[\phi(j) - \phi(\alpha)]} V_{\alpha j}$$

where $\alpha(j)$ denotes and up-kind (down-kind) quark.

This rephasing of L-fields only affects the term $\mathcal{L}^{phys}(f, H)$ in the Lagrangian density, but it can be remedied by rephasing the R-fields, so that the Lagrangian density remains invariant. A general unitary V matrix has N^2 parameters, with

$$\frac{N(N-1)}{2} \text{ moduli} \quad , \quad \frac{N(N+1)}{2} \text{ phases}$$

From rephasing invariance, we can absorb $(2N-1)$ unobservable phases, so that¹⁴ one finishes with $(N-1)^2$ physical parameters, among which

$$\frac{N(N-1)}{2} \text{ moduli} \quad , \quad \frac{(N-1)(N-2)}{2} \text{ phases}$$

For the case of 2 families, $N = 2$, we find 1 rotation angle and no phases. The quark mixing matrix is then the Cabibbo matrix¹⁵, which is real orthogonal. Reality implies that CP is conserved in the Standard Model with 2 families.

For 3 families, $N = 3$, the number of physical parameters in V is 4, i.e., 3 rotation angles and 1 phase. The first example of such a matrix was given by Kobayashi and Maskawa¹⁶.

In order to have CP-violation, the three families have to be active in the process at hand. If two quarks with the same charge were degenerate in mass, there is an extra symmetry

$$(s, b) \rightarrow s^{new} = V_{us}s + V_{ub}b$$

such that u is not coupled to b^{new} , and the new V could be chosen real. Necessary conditions for CP-violation are

$$\begin{aligned} m_u &\neq m_c & , & & m_c &\neq m_t & , & & m_t &\neq m_u \\ m_d &\neq m_s & , & & m_s &\neq m_b & , & & m_b &\neq m_d \end{aligned}$$

There are unfortunately many possible parametrizations of V . PDG advocates¹⁷ the following form

$$V = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{13}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{13}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{13}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{13}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{13}} & c_{23}c_{13} \end{bmatrix}$$

where the rotation angles $\theta_{12}, \theta_{23}, \theta_{13}$ can be made to lie in the first quadrant, so that

$$c_{ij} \geq 0 \quad , \quad s_{ij} \geq 0 \quad ; \quad 0 \leq \delta_{13} \leq 2\pi$$

A very interesting approximate parametrization is due to Wolfenstein¹⁸, in terms of the four parameters

$$(\lambda, A, \rho, \eta)$$

with λ , the Cabibbo angle, taken as the expansion parameter:

$$V = \begin{bmatrix} 1 - \frac{1}{2}\lambda^2 & \lambda & \lambda^3 A(\rho - i\eta) \\ -\lambda & 1 - \frac{1}{2}\lambda^2 & \lambda^2 A \\ \lambda^3 A(1 - \rho - i\eta) & -\lambda^2 A & 1 \end{bmatrix}$$

In this parametrization, CP-violation is manifested in the presence of an imaginary part of the V_{ub}, V_{td} matrix elements: $\eta \neq 0$.

2 The $K^0 - \bar{K}^0$ system

2.1 Discovery of CP-Violation

One consequence of CP-INVARIANCE for the neutral $K^0 - \bar{K}^0$ system was predicted by Gell-Mann and Pais¹⁹: there should be a long-lived partner to the known $V^0(K_1^0)$ particle of short lifetime: $\tau(K_s) \sim 10^{-10}$ sec. According to this proposal these two particles are coherent superpositions of two strangeness eigenstates, $K^0(S = +1)$ and $\bar{K}^0(S = -1)$, produced in interactions with conservation of strangeness, like strong interactions. Weak interactions do not conserve strangeness and the physical particles should be eigenstates of CP if the weak interactions are CP-invariant. These eigenstates are (with $\bar{K}^0 \equiv -CP K^0$):

$$CP K_1 = CP \frac{1}{\sqrt{2}}(K^0 - \bar{K}^0) = \frac{1}{\sqrt{2}}(-\bar{K}^0 + K^0) = K_1$$

$$CP K_2 = CP \frac{1}{\sqrt{2}}(K^0 + \bar{K}^0) = \frac{1}{\sqrt{2}}(-\bar{K}^0 - K^0) = -K_2$$

K_1, K_2 , eigenstates of CP, should be the physical particles with definite mass and lifetime.

Because of $CP(\pi^+\pi^-) = (\pi^+\pi^-)$, $CP(\pi^0\pi^0) = (\pi^0\pi^0)$, the decay into $(\pi\pi)$ is allowed for K_1 , to be identified with the experimental state K_s of short lifetime. The decay into $(\pi\pi)$ is forbidden for K_2 , hence its longer lifetime, K_L , which was indeed confirmed when discovered²⁰.

In 1964, Christenson, Cronin, Fitch and Turlay⁹ discovered that K_L also decays to $\pi^+\pi^-$ with a branching ratio of $\sim 2 \times 10^{-3}$. This discovery established CP-VIOLATION.

Then, the long-lived neutral Kaon K_L is no longer identical to the CP-eigenstate K_2 . Similarly, K_s is not identical to K_1 . CP-Violation, discovered by the decay $K_L \rightarrow \pi^+\pi^-$, was confirmed later by $K_L \rightarrow \pi^0\pi^0$ ²¹ and by the charge asymmetry²² in the K_{l3} decays: $K_L \rightarrow \pi^\pm e^\mp \nu$ and $K_L \rightarrow \pi^\pm \mu^\mp \nu$.

Early models for CP-violation attributed this phenomenon to a fifth force, the superweak interaction²³, to a T-odd part of the weak interaction or to

the interference with a T-odd part of the electromagnetic or strong interaction. Many subsequent experiments have reduced the possible models to the superweak and milliweak classes. Amongst the milliweak models, the Present Orthodoxy is based on the Complex Quark Mixing Matrix for 3 families

2.2 Meson-Antimeson Mixing

Let the eigenstates of flavour be $P^0(=K^0, D^0, B^0)$ and $\bar{P}^0$. Flavour Number is not conserved by H_w of weak interactions, so in 2^{nd} order P^0 and $\bar{P}^0$ can mix through the intermediate states connected by H_w . The mixing leads to

$$|\psi(t)\rangle = a(t)|P^0\rangle + b(t)|\bar{P}^0\rangle \equiv \begin{Bmatrix} a(t) \\ b(t) \end{Bmatrix}$$

with the Time Evolution governed by the Mixing Matrix $\mathcal{M}$. Assuming CPT-invariance,

$$i\frac{d}{dt}|\psi(t)\rangle = \mathcal{M}|\psi(t)\rangle$$

$$\mathcal{M} = \begin{pmatrix} M & M_{12} \\ M_{12}^* & M \end{pmatrix} - \frac{i}{2} \begin{pmatrix} \Gamma & \Gamma_{12} \\ \Gamma_{12}^* & \Gamma \end{pmatrix}$$

The non-hermiticity of $\mathcal{M}$ is associated with the existence of decay channels, not introduced explicitly in the formalism. The values of M and Γ , real, would correspond to the mass and width of both $P^0, \bar{P}^0$ in absence of Mixing.

The off-diagonal entries M_{12}, Γ_{12} of the mass matrix $\mathcal{M}$ correspond to the Dispersive and Absorptive parts, respectively, of the $\Delta P = 2$ transitions. If CP were a good symmetry, then M_{12}, Γ_{12} would be real.

The physical eigenstates of $\mathcal{M}$ are

$$|P_{\mp}\rangle = \frac{1}{\sqrt{|p|^2 + |q|^2}} [p|P^0\rangle \mp q|\bar{P}^0\rangle]$$

with the amplitudes

$$\frac{q}{p} \equiv \frac{1 - \bar{\varepsilon}}{1 + \bar{\varepsilon}} = \left(\frac{M_{12}^* - \frac{i}{2}\Gamma_{12}^*}{M_{12} - \frac{i}{2}\Gamma_{12}} \right)^{1/2}$$

In the limit of CP-symmetry, M_{12} and Γ_{12} are real and thus $q/p = 1$: the physical eigenstates become eigenstates of CP

$$|P_{1,2}\rangle = \frac{1}{\sqrt{2}} (|P^0\rangle \mp |\bar{P}^0\rangle) ; CP|P_{1,2}\rangle = \pm |P_{1,2}\rangle$$

In the presence of CP-violation, the physical eigenstates $|P_{\pm}\rangle$ are NOT orthogonal, a result implied by the non-hermiticity of $\mathcal{M}$. One has

$$\langle P_- | P_+ \rangle = \frac{|p|^2 - |q|^2}{|p|^2 + |q|^2} \approx 2 \operatorname{Re} \bar{\varepsilon}$$

With the initial condition ($t = 0$) that the produced state is either P^0 or $\bar{P}^0$, the corresponding Time Evolution gives

$$\begin{Bmatrix} |P^0(t)\rangle \\ |\bar{P}^0(t)\rangle \end{Bmatrix} = \begin{bmatrix} g_1(t) & \frac{q}{p}g_2(t) \\ \frac{p}{q}g_2(t) & g_1(t) \end{bmatrix} \begin{Bmatrix} |P^0\rangle \\ |\bar{P}^0\rangle \end{Bmatrix}$$

where the two time-dependent functions $g_{1,2}(t)$ are

$$\begin{Bmatrix} g_1(t) \\ g_2(t) \end{Bmatrix} = e^{-iMt} e^{-\Gamma t/2} \begin{Bmatrix} \cos(\Delta M - \frac{i}{2}\Delta\Gamma)\frac{t}{2} \\ -i \sin(\Delta M - \frac{i}{2}\Delta\Gamma)\frac{t}{2} \end{Bmatrix}$$

with $\Delta M \equiv M_{P_+} - M_{P_-}$; $\Delta\Gamma \equiv \Gamma_{P_+} - \Gamma_{P_-}$

The main difference between the $K^0 - \bar{K}^0$ system and the $B^0 - \bar{B}^0$ system lies in the relative values of these parameters:

$$\begin{aligned} -K \text{ system, } & \Gamma_{K_L} \ll \Gamma_{K_S}, \quad \Delta\Gamma_K \approx -\Gamma_{K_S} \approx -2\Delta M_K \\ -B \text{ system, } & \Delta\Gamma_B \ll \Gamma_B, \quad \Delta\Gamma_B \ll \Delta M_B \end{aligned}$$

2.3 Indirect CP-Violation

The flavour-specific decays

$$\begin{aligned} K^0 & \rightarrow \pi^- l^+ \nu_l \\ \bar{K}^0 & \rightarrow \pi^+ l^- \bar{\nu}_l \end{aligned}$$

can measure whether $|p/q| \neq 1$, i.e., CP-violation in the physical eigenstates of the Mass Matrix, referred to as Indirect CP-violation.

In the Standard Model, the semileptonic decay amplitudes from K^0 or $\bar{K}^0$ are equal

$$|A(\bar{K}^0 \rightarrow \pi^+ l^- \bar{\nu}_l)| = |A(K^0 \rightarrow \pi^- l^+ \nu_l)|$$

so that the charge asymmetry from K_L -decays is given by

$$\delta = \frac{\Gamma(K_L \rightarrow \pi^- l^+ \nu_l) - \Gamma(K_L \rightarrow \pi^+ l^- \bar{\nu}_l)}{\Gamma(K_L \rightarrow \pi^- l^+ \nu_l) + \Gamma(K_L \rightarrow \pi^+ l^- \bar{\nu}_l)} = \frac{|p|^2 - |q|^2}{|p|^2 + |q|^2}$$

and indicates that all the CP-violating effect comes from the selection of either K^0 or $\bar{K}^0$ in the K_L -state.

Present values of δ give a world average¹⁷

$$\delta = (3.27 \pm 0.12) \times 10^{-3}$$

so that there is

CP-Violation in the Mixing
 $Re \bar{\varepsilon}_K = (1.63 \pm 0.06) \times 10^{-3}$

2.4 Isospin Decomposition

For the non-leptonic $K_{S,L} \rightarrow 2\pi$ decays, the angular momentum of the pions vanishes. The spatial part of the (2π) -state is therefore symmetric, and since pions are bosons, the isospin state must be symmetric too. The two symmetric combinations of two $I = 1$ states have $I = 0$ and $I = 2$, and the four existing transition amplitudes are

$$\langle 0|T|K_s \rangle \quad , \quad \langle 2|T|K_s \rangle \quad , \quad \langle 0|T|K_L \rangle \quad , \quad \langle 2|T|K_L \rangle$$

By normalizing to the dominant CP-conserving $\Delta I = 1/2$ transition amplitude $\langle 0|T|K_s \rangle$, we define three complex numbers

$$\begin{aligned} \varepsilon_0 &\equiv \langle 0|T|K_L \rangle / \langle 0|T|K_s \rangle \\ \varepsilon_2 &\equiv \frac{1}{\sqrt{2}} \langle 2|T|K_L \rangle / \langle 0|T|K_s \rangle \\ \omega &\equiv \langle 2|T|K_s \rangle / \langle 0|T|K_s \rangle \end{aligned}$$

Non-vanishing values of $\varepsilon_{0,2}$ are a demonstration of CP-violation, whereas ω parametrizes the relative contribution of the $\Delta I = 3/2$ amplitude.

The experimentally observable quantities are

$$\begin{aligned} \eta_{+-} &= \langle \pi^+ \pi^- | T | K_L \rangle / \langle \pi^+ \pi^- | T | K_s \rangle \\ \eta_{00} &= \langle \pi^0 \pi^0 | T | K_L \rangle / \langle \pi^0 \pi^0 | T | K_s \rangle \end{aligned}$$

Relating the isospin states to the physical (2π) -states,

$$\begin{aligned} \langle 0| &= \frac{1}{\sqrt{3}} \langle \pi^- \pi^+ | - \frac{1}{\sqrt{3}} \langle \pi^0 \pi^0 | + \frac{1}{\sqrt{3}} \langle \pi^+ \pi^- | \\ \langle 2| &= \frac{1}{\sqrt{6}} \langle \pi^- \pi^+ | + \sqrt{\frac{2}{3}} \langle \pi^0 \pi^0 | + \frac{1}{\sqrt{6}} \langle \pi^+ \pi^- | \end{aligned}$$

one obtains

$$\eta_{+-} = \frac{\varepsilon_0 + \varepsilon_2}{1 + \frac{\omega}{\sqrt{2}}} \quad , \quad \eta_{00} = \frac{\varepsilon_0 - 2\varepsilon_2}{1 - \sqrt{2}\omega}$$

Because of the validity of the $\Delta I = 1/2$ rule for CP-conserving weak non-leptonic decays, one expects, $|\omega| \ll 1$. Its actual value gives $|\omega| \sim 1/22$.

A suitable choice for the phase of the $K^0 \rightarrow 2\pi$ ($I = 0$) amplitude is obtained by putting all its complexity as coming from final-state-interactions leading to a phase-shift δ_0 :

$$\left. \begin{aligned} \langle 0|T|K^0 \rangle &\equiv iA_0 e^{i\delta_0} \\ \langle 0|T|\bar{K}^0 \rangle &\equiv -iA_0^* e^{i\delta_0} \end{aligned} \right\} \quad \begin{aligned} A_0 &\text{real,} \\ \text{Im}(A_0) &= 0 \end{aligned}$$

Similarly, one has

$$\left. \begin{aligned} \langle 2|T|K^0 \rangle &\equiv iA_2 e^{i\delta_2} \\ \langle 2|T|\bar{K}^0 \rangle &\equiv -iA_2^* e^{i\delta_2} \end{aligned} \right\}$$

Then one gets

$$\varepsilon_0 = \varepsilon_K \quad , \quad \varepsilon_2 = i \frac{1}{\sqrt{2}} \frac{\text{Im} A_2}{A_0} e^{i(\delta_2 - \delta_0)} \equiv \varepsilon'$$

as well as the charge asymmetry $\delta = 2 \text{Re} \varepsilon_K$

We observe:

- (i) ε_K measures indirect CP-violation
- (ii) ε' measures direct CP-violation, in the decay amplitude, governed by the phase-shift in final-state-interaction¹⁷ $\delta_2 - \delta_0 = -45^\circ \pm 6^\circ$
- (iii) The observable quantities satisfy the Wu-Yang triangle relations

$$\eta_{+-} = \varepsilon + \varepsilon' \quad , \quad \eta_{00} = \varepsilon - 2\varepsilon'$$

The experimental objective is the separation of ε and ε'

The phase of ε' is $\text{Arg}(\varepsilon') = \frac{\pi}{2} + (\delta_2 - \delta_0) = (45^\circ \pm 6^\circ)$.

The phase of ε can be identified from the diagonalization of the mass matrix $\mathcal{M}$.

$$\varepsilon_K \simeq e^{i\phi_{sw}} \frac{\text{Im} M_{12} - \frac{i}{2} \text{Im} \Gamma_{12}}{\sqrt{\Delta M_K^2 + \frac{1}{4} \Delta \Gamma_K^2}}$$

where ϕ_{sw} is the so-called superweak phase

$$\phi_{sw} \equiv \arg \tan\left(\frac{-2\Delta M_K}{\Delta \Gamma_K}\right)$$

Taking into account $\Delta \Gamma_K \simeq -2\Delta M_K$, we discover that ϕ_{sw} is again near to $\pi/4$. More precisely, $\phi_{sw} = 43.64^\circ \pm 0.15^\circ$.

The phase introduced by the second factor in ε_K is very small, so one concludes that the phases of ε and ε' are nearly equal. The equality of the phases ϕ_{+-} and ϕ_{00} of the observable amplitudes η_{+-} and η_{00} , respectively, is a test of CPT-invariance in Nature.

2.5 Experiments for $K_L \rightarrow 2\pi$

The experimental results

$$|\eta_{+-}| = (2.269 \pm 0.023) \times 10^{-3}, \quad |\eta_{00}| = (2.259 \pm 0.023) \times 10^{-3}$$

show their equality within errors, indicating that $|\varepsilon'| \ll |\varepsilon|$. This is a consequence of the approximate $\Delta I = 1/2$ rule.

In order to extract the corresponding phases ϕ_{+-}, ϕ_{00} , one needs a mechanism of $K_L - K_s$ interference. This is possible from either (i) Coherent Regeneration of K_s from K_L , or (ii) The Time-Distribution of $K \rightarrow \pi\pi$ events as obtained from a pure flavour (strangeness) state prepared at $t = 0$.

The first method (i) needs the preparation of K_L -beams. The second method (ii) makes use of flavour tag: either K^0 or $\bar{K}^0$. From the results on $K_L \rightarrow 2\pi$, one reproduces $Re \varepsilon_K \simeq 1.63 \times 10^{-3}$, in good agreement with the value extracted from the charge asymmetry δ in semileptonic decays. The second method (ii) has been put into practice by the CP-LEAR Collaboration at CERN recently²⁴, and their results for $\pi^+\pi^-$ Decays are shown in Figure 2.


These results represent the first direct observation of a difference in the decay rates between particles and antiparticles.

The ratio ε'/ε can be determined by the "method of ratio of ratios" when comparing the $\pi^0\pi^0$ and $\pi^+\pi^-$ decay channels:

$$Re(\frac{\varepsilon'}{\varepsilon}) \simeq \frac{1}{6} \left\{ 1 - \left| \frac{\eta_{00}}{\eta_{+-}} \right|^2 \right\}$$

Present results give $(23.0 \pm 6.5) \times 10^{-4}$ for the NA31- Collaboration²⁵ at CERN and $(7.4 \pm 5.9) \times 10^{-4}$ for the E731-Collaboration²⁶ at FermiLab. These values are statistically compatible within 8%. New experiments with better sensitivity at CERN, FermiLab and a dedicated Φ -Factory at Frascati will push the precision to better values. The Goal is to reach sensitivities better than 10^{-4} !

CP Violation in $\pi^+ \pi^-$ Decays



First direct observation of difference of decay rates between particles and antiparticles.

Figure 2: Time Evolution of K^0 and $\bar{K}^0$ rates at CP-LEAR

2.6 CKM Quark Mixing Matrix

The standard mechanism to incorporate CP-violation in the present electroweak theory is the Cabibbo-Kobayashi-Maskawa Complex Mixing Matrix for 3 families of Quarks. It generates CP-violating effects in both: (i) the $\Delta S = 2$ $K^0 \bar{K}^0$ transition, through the

Box-Diagram




Figure 3: Quark diagram for $\Delta S = 2$ $K^0 \rightleftharpoons \bar{K}^0$

and (ii) the $\Delta S = 1$ Decay amplitude to non-strange quarks, through the

Penguin-Diagram




Figure 4: Quark diagram for $\Delta S = 1$ K^0 Decays

The real part of the $K^0 - \bar{K}^0$ mixing, which contributes to ΔM , is dominated by intermediate charm quarks running in the loop. Its main uncertainty

comes from the hadronic matrix element of the effective four-quark $\Delta S = 2$ operator, when the higher degrees of freedom are integrated out. Its value is parametrized by B_K , measuring it with respect to the vacuum-contribution in the intermediate state.

The imaginary part of the box-diagram provides the explanation of the indirect CP-violation ε_K in the Standard Model and it is dominated by intermediate top quarks running in the loop. Its experimental value can be analyzed in terms of the V_{td} matrix-element of the CKM Mixing Matrix.

Using the Wolfenstein parametrization, the experimental value of ε_K specifies a hyperbola in the (ρ, η) plane: the allowed region is shown in Fig. 5.




Figure 5: Allowed region in the (ρ, η) plane, constrained from the experimental value of ε_K .

A theoretical estimate of the direct CP-violation ε'/ε is more involved. The quark diagrams giving $\Delta S = 1$ transitions, including the penguins, induce up to 10 four-quark operators once the higher degrees of freedom are integrated out. A control of the long-distance effects becomes both fundamental and difficult. Although the gluon-mediated penguin was initially assumed to be the main ingredient, the Z -penguin contains non-decoupling m_t^2 terms in the amplitude. The final result is thus very sensitive to m_t . The present theory provides values for ε'/ε ranging from -3×10^{-4} to 10^{-3} , depending on the hadronic physics being taken by the Roma, München or Dortmund groups.

2.7 Coherent Decay of $(K^0 \bar{K}^0)$

The two-particle $K^0 - \bar{K}^0$ system, such as is being produced by Φ -decay, has a total angular momentum $J = L$ and it is eigenstate of charge-conjugation C with eigenvalue $C = (-1)^L$.

Bose statistics requires that the physical state has to be symmetric under the product of $C \times \mathcal{P}$, where C is charge-conjugation and $\mathcal{P}$ is permutation of space coordinates. In other words, the physical state $|\psi\rangle$ has to be eigenstate of $C\mathcal{P}$ with eigenvalue $+1$.

Thus, as a consequence of Quantum Mechanics indistinguishability, we can write

$$\begin{aligned} |\psi\rangle &= \frac{1}{\sqrt{2}}(1 + C\mathcal{P})|LM; K^0 \bar{K}^0\rangle \\ &= \frac{1}{\sqrt{2}} \left\{ |K^0(\vec{k}) \bar{K}^0(-\vec{k})\rangle + (-1)^L |\bar{K}^0(\vec{k}) K^0(-\vec{k})\rangle \right\} \end{aligned}$$

If we impose the conditions of the e^+e^- machine at the Φ -peak (like DAΦNE at Frascati), we have $L = 1$ and $C = -$. The produced state of $(K^0 - \bar{K}^0)$, when written in terms of the physical neutral kaon states with definite mass and lifetime, is

$$|K^0 \bar{K}^0(C = -)\rangle = \frac{1 + |\alpha|^2}{2\sqrt{2}\alpha} \left\{ |K_L(\vec{k}) K_s(-\vec{k})\rangle - |K_s(\vec{k}) K_L(-\vec{k})\rangle \right\}$$

with no component of $K_s K_s$ or $K_L K_L$ states.

We conclude²⁷ :

- (i) Bose statistics for $L = 1$ says that the state of two identical bosons is forbidden !
- (ii) The result (i) applies to any time in the evolution of the system, and thus we have simultaneously NO $K^0 K^0$, NO $\bar{K}^0 \bar{K}^0$!
- (iii) The preparation of $(K^0 \bar{K}^0)$ from a Φ -factory leads to correlated (simultaneous) beams of K_L and K_s !

The parameter α in the normalization of the state is the ratio of the amplitudes

$$\alpha \equiv \frac{p}{q} = \frac{1 + \varepsilon_K}{1 - \varepsilon_K}$$

By studying the decay amplitude from this correlated system, we ask for the appearance of X_1 at time t_1 and X_2 at time t_2 , as illustrated in Figure 6




Figure 6: The Decay of the Correlated ($K^0 \bar{K}^0$)

$$A(X_1, X_2) = \frac{1+|\alpha|^2}{2\sqrt{2}\alpha} < X_1 | K_s > < X_2 | K_s > e^{-i(\lambda_L + \lambda_s)t/2} \\ \{ \eta_1 e^{-i\Delta\lambda\Delta t/2} - \eta_2 e^{i\Delta\lambda\Delta t/2} \}$$

where

$$\eta_j = \frac{< X_j | K_L >}{< X_j | K_s >} \quad ; \quad j = 1, 2$$

and

$$\lambda_{s,L} = M_{s,L} - \frac{i}{2}\Gamma_{s,L} \quad ; \quad \Delta\lambda = \lambda_s - \lambda_L ; \\ t = t_1 + t_2 \quad ; \quad \Delta t = t_2 - t_1$$

Suppose that we detect the two decay products X_1 and X_2 at equal times $t_1 = t_2$. For $\Delta t = 0$, the decay amplitude gives

$$A(X_1, X_2; \Delta t = 0) \propto (\eta_1 - \eta_2)$$

In particular, for $X_1 = X_2$ one finds

$$A[X_1(t_1), X_1(t_1)] = 0$$

which is a quantum mechanical correlation²⁸ of the Einstein-Podolsky-Rosen type implied by Bose statistics.

If we select the two different 2π decay channels of the neutral kaon, i.e. $X_1 \equiv \pi^+\pi^-$ and $X_2 \equiv \pi^0\pi^0$, we find at equal times

$$A[\pi^+\pi^-(t_1), \pi^0\pi^0(t_1)] \propto (\eta_{+-} - \eta_{00}) = 3\varepsilon'$$

This result has been discussed²⁹ in the context of Φ -factories, trying to maximize the ε' effect by comparing the observables at different time slices. The intensity asymmetry, for a given Δt , under the change $\Delta t \rightarrow -\Delta t$, is very sensitive to ε'/ε when Δt is of the order of τ_s , the K_s lifetime.

2.8 Time Integrated Rates

Intead of comparing the rates at different time intervals, for a given decay channel $\pi^+\pi^-\pi^0\pi^0$ of the $K^0 - \bar{K}^0$ system, one can adopt a different strategy: the comparison of different decay chanelns in time integrated rates³⁰.

In particular,

$$\frac{Br(\pi^+\pi^-, \pi^+\pi^-)}{Br(\pi^0\pi^0, \pi^0\pi^0)} = \left[\frac{Br(K_s \rightarrow \pi^+\pi^-)}{Br(K_s \rightarrow \pi^0\pi^0)} \right]^2 \left\{ 1 + 6Re\left(\frac{\varepsilon'}{\varepsilon}\right) \right\}$$

and similarly³⁰ for the other possible ratios.

3 CP- Violation and B-Physics

3.1 Physics Motivation

The main question is whether the origin of CP-violation can be explained within the Standard Model or it needs physics beyond the Standard Model. Inside the Standard Model, the analysis for Flavour Mixing starts from a Charged Current Lagrangian, with the Quark Mixing Matrix described by 4 independent parameters. In the Wolfenstein parametrization, for example, these are $[\lambda, A, \rho, \eta]$.

It has to be emphasized that these are fundamental couplings of the Standard Model Lagrangian, so their determination is of primordial importance. In order to answer the question on the possible need of beyond-the-standard-model physics, one has to look for ways to overdetermine these parameters and show the internal consistency of the model.

From the Unitary Quark Mixing Matrix

$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{bmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{bmatrix}$$

one can establish 6 off-diagonal Unitarity Relations which, represented in the complex plane, can be visualized (for 3 families) by 6 Unitarity Triangles. One of these, that built from the d - and b - columns, is written as

$$(b, d) \rightarrow V_{ub}^* V_{ud} + V_{cb}^* V_{cd} + V_{tb}^* V_{td} = 0$$

and it has the peculiarity that the three sides are of the same order of magnitude: $O(\lambda^3)$.

In the Standard Model, a single phase parameter (η) is the only possible source of CP violation, so that the predictions for CP-violating phenomena are quite constrained. Moreover, the CKM mechanism requires several necessary conditions in order to generate an observable CP-violating effect: (i) All three families are required to play an active role. With only two fermion families, the Quark Mixing mechanism cannot give rise to CP-violation. In the kaon system, for instance, CP-violation effects can only appear at the one-loop level, where the top quark is present. In B-physics, however, the decays generated by tree level diagrams can induce CP-violation effects; (ii) The quarks of a given charge must be non-degenerate in mass. With degeneracy, the physical quark states could be redefined in order to vanish the quark mixing matrix element; (iii) all CKM-matrix elements must be non-zero. If any of these conditions were not satisfied, the CKM-phase could be rotated away by a redefinition of the quark fields. CP-violation effects are then necessarily proportional to the product of all CKM angles.

All these necessary conditions are summarized in a single requirement³¹ on the original quark-mass matrices M_u and M_d :

$$CP - \text{violation} \iff \text{Im}\{\det[M_u M_u^\dagger, M_d M_d^\dagger]\} \neq 0$$

In the case of 3 families, there is a unique combination of angles and phases to generate CP-violation for all flavours that can be considered:

$$J \simeq A^2 \lambda^6 \eta \leq 10^{-4}$$

Any CP-violation observable involves the product J and, thus, violations of the CP symmetry are necessarily small.

One can make some general statements:

(i) In order to generate sizeable CP-violating asymmetries $[(\Gamma - \bar{\Gamma})/(\Gamma + \bar{\Gamma})]$ one should look for very suppressed decays, where the decay widths already involve small CKM matrix elements.

(ii) In the Standard Model, CP violation is a low-energy phenomenon, in the sense that any effect should disappear with $\frac{m_c^2 - m_u^2}{s} \ll 1$.

(iii) B decays are the optimal place for CP-violation signals to show up. They involve small CKM matrix elements in the decays and are the lowest mass processes where the three quark families play a tree-level role.

The (bd) Unitarity Triangle is shown in Fig. 7, where it has been scaled by dividing its sides by $|V_{cb}^* V_{cd}|$. In the Wolfenstein parametrization, this is real. Aligning it, with length equal to 1, along the real axis, the coordinates of the 3 vertices are then $(0,0)$, $(1,0)$ and (ρ, η) . Note that, although the orientation of the triangle in the complex plane is phase-convention dependent, the triangle

itself is a physical object: the length of the sides and the angles can be directly measured.




Figure 7: The (bd) Unitarity Triangle

The sides of the Unitarity Triangle are determined from the measured ratio $\Gamma(b \rightarrow u)/\Gamma(b \rightarrow c)$ and from $B_d^0 - \bar{B}_d^0$ mixing:

$$R_b = \left| \frac{V_{ub}^* V_{ud}}{V_{cb}^* V_{cd}} \right| \approx \left| \frac{V_{ub}}{\lambda V_{cb}} \right| \approx \sqrt{\rho^2 + \eta^2}$$

$$R_t = \left| \frac{V_{tb}^* V_{td}}{V_{cb}^* V_{cd}} \right| \approx \left| \frac{V_{td}}{\lambda V_{cb}} \right| \approx \sqrt{(1 - \rho)^2 + \eta^2}$$

The significance of the V_{td} coupling is seen in the box diagram of Figure 8, relevant to the $B_d^0 - \bar{B}_d^0$ mixing. A priori, the measurement of the two sides, performed through CP-conserving observables, could make possible to establish that the area is not vanishing and that Fig. 7 is indeed a triangle. With the present theoretical and experimental uncertainties, this is however not possible.




Figure 8: Quark diagram responsible of the $B_d^0 - \bar{B}_d^0$ mixing. The arrows indicate the V_{td} coupling.

In section 2 we have seen that the experimental value of ϵ_K specifies a hyperbola in the (ρ, η) plane. This constraint, together with the constraints from

R_b and R_t , which result in the circles centered at $(0, 0)$ and $(1, 0)$, respectively, give an allowed range of values³² for (ρ, η) as shown in Fig. 9. The final value is given by the intersection of all constraints.




Figure 9: Constraints for (ρ, η) coming from the measured values of R_b, R_t and ϵ_K .

3.2 Principle of the CP-violation measurement

i) With a large $B^0 - \bar{B}^0$ mixing we can generate a CP-violating interference in the following way. There are quite a few non-leptonic final states which are reachable both from a B^0 and a $\bar{B}^0$. For these flavour non-specific decays the B^0 (or $\bar{B}^0$) can decay directly to a given final state f , or do it after the meson has been changed to its antiparticle via the mixing process; i.e., there are two different amplitudes, $A(B^0 \rightarrow f)$ and $A(B^0 \rightarrow \bar{B}^0 \rightarrow f)$, corresponding to two possible decay paths, as shown in Fig. 10.




Figure 10: Two decay paths from B^0 to f , with interfering amplitudes.

CP-violating effects can then result³³ from the interference of these two am-

plitudes. To build the associated asymmetry one needs to TAG the initial B-flavour.

The time-dependent decay probabilities for the decay of a neutral B meson created at the time $t_o = 0$ as a pure $B^0(\bar{B}^0)$ into the final state $f(\bar{f} \equiv CP f)$ are (we neglect the tiny $\Delta\Gamma_{B^0}$ corrections):

$$\begin{aligned}\Gamma[B^0(t) \rightarrow f] &\sim \frac{1}{2}e^{-\Gamma t}|A_f|^2 \left\{ [1 + |\bar{\rho}_f|^2] + \right. \\ &\quad \left. + [1 - |\bar{\rho}_f|^2] \cos(\Delta Mt) - 2\text{Im}(\frac{q}{p}\bar{\rho}_f) \sin(\Delta Mt) \right\} \\ \Gamma[\bar{B}^0(t) \rightarrow \bar{f}] &\sim \frac{1}{2}e^{-\Gamma t}|\bar{A}_f|^2 \left\{ [1 + |\rho_f|^2] + \right. \\ &\quad \left. + [1 - |\rho_f|^2] \cos(\Delta Mt) - 2\text{Im}(\frac{p}{q}\rho_f) \sin(\Delta Mt) \right\}\end{aligned}$$

where we have introduced the notation

$$\begin{aligned}A_f &\equiv A[B^0 \rightarrow f], & \bar{A}_f &\equiv -A[\bar{B}^0 \rightarrow f], & \bar{\rho}_f &\equiv \bar{A}_f/A_f \\ A_{\bar{f}} &\equiv A[B^0 \rightarrow \bar{f}], & \bar{A}_{\bar{f}} &\equiv -A[\bar{B}^0 \rightarrow \bar{f}], & \rho_{\bar{f}} &\equiv A_{\bar{f}}/\bar{A}_{\bar{f}}\end{aligned}$$

CP-invariance demands the probabilities of CP conjugate processes to be identical. Thus, CP conservation requires $A_f = \bar{A}_{\bar{f}}, A_{\bar{f}} = \bar{A}_f, \bar{\rho}_f = \rho_{\bar{f}}$ and $\text{Im}(\frac{q}{p}\bar{\rho}_f) = \text{Im}(\frac{p}{q}\rho_{\bar{f}})$. The violation of any of the first three equalities would be a signal of direct CP-violation, i.e., in the decay amplitudes. The fourth equality tests CP violation generated by the interference of the direct decay $B^0 \rightarrow f$ and the mixing-induced decay $B^0 \rightarrow \bar{B}^0 \rightarrow f$, as shown in Fig. 10.

If the final state is a CP-eigenstate, $\bar{f} = \tau_f f$ with $\tau_f = \pm 1$ and there is only one transition decay amplitude^b, we have

$$\rho_{\bar{f}} = \bar{\rho}_f = \tau_f e^{2i\phi_{WD}}$$

$$\frac{\Gamma[B^0(t) \rightarrow f] - \Gamma[\bar{B}^0(t) \rightarrow f]}{\Gamma[B^0(t) \rightarrow f] + \Gamma[\bar{B}^0(t) \rightarrow f]} = \tau_f \sin(2\Phi) \sin(\Delta Mt)$$

where Φ is the convention-independent CP-phase which results from the combination of the phase ϕ_{WD} of the decay amplitude and the phase of the mixing parameter $\frac{q}{p}$. This mechanism is thus called "the interplay between Mixing and Direct CP-Violation".

ii) Consider now two interfering decay amplitudes which contribute to a given process $B \rightarrow f$:

$$\begin{aligned}A(B \rightarrow f) &= M_1 e^{i\phi_1} e^{i\alpha_1} + M_2 e^{i\phi_2} e^{i\alpha_2} \\ A(\bar{B} \rightarrow \bar{f}) &= M_1 e^{-i\phi_1} e^{i\alpha_1} + M_2 e^{-i\phi_2} e^{i\alpha_2}\end{aligned}$$

^bEach transition amplitude is identified as the weak interaction matrix element of a given operator with a definite CP-phase

where ϕ_1, ϕ_2 are the corresponding weak decay phases and α_1, α_2 are the final state "strong" interaction phases. They are such that, in going to the CP-conjugated process, the weak phases change their sign, whereas the strong phases keep their sign. The strong phases originate from the beyond the Born approximation complexity of the decay amplitudes. The rate Asymmetry is thus given by ³⁴

$$\frac{\Gamma(B \rightarrow f) - \Gamma(\bar{B} \rightarrow \bar{f})}{\Gamma(B \rightarrow f) + \Gamma(\bar{B} \rightarrow \bar{f})} = \frac{-2M_1M_2 \sin(\phi_1 - \phi_2) \sin(\alpha_1 - \alpha_2)}{|M_1|^2 + |M_2|^2 + 2M_1M_2 \cos(\phi_1 - \phi_2) \cos(\alpha_1 - \alpha_2)}$$

so that one needs ³⁴:

- 1) Two, at least, interfering amplitudes.
- 2) Two different weak phases $\phi_1 \neq \phi_2$
- 3) Two different strong phases $\alpha_1 \neq \alpha_2$.
- 4) To enhance the asymmetry, M_1 and M_2 should be of comparable size.

This "Direct CP-Violation" operates either for charged B's or neutral B's with flavour specific decays.

iii) There is a third method to generate CP-Violation, that of CP-Violation in $B^0 - \bar{B}^0$ Mixing or "Indirect CP-Violation". Take the semileptonic decays of a $B^0 - \bar{B}^0$ pair with equal sign dileptons: $N(l^+l^+)$ or $N(l^-l^-)$ is a signal of mixing, their difference is a signal of CP-Violation in this mixing.

One gets the semileptonic asymmetry of $B^0 \rightarrow Xl\nu_l$

$$a_{SL} = \frac{N(l^+l^+) - N(l^-l^-)}{N(l^+l^+) + N(l^-l^-)} = \frac{|p/q|^2 - |q/p|^2}{|p/q|^2 + |q/p|^2} \simeq 4Re \bar{\epsilon}_B$$

The off-diagonal matrix elements of the effective mixing matrix of $B^0 - \bar{B}^0$ give rise to a difference of the masses and widths of the two physical eigenstates. In the Standard Model the two complex eigenvalues satisfy

$$\left| \frac{\Delta\Gamma_B}{\Delta M_B} \right| \sim \left| \frac{\Gamma_{12}}{M_{12}} \right| \sim \frac{m_b^2}{m_t^2}$$

To generate $Re(\bar{\epsilon}_B) \neq 0$ one needs both $\Delta\Gamma_B \neq 0$ and a misalignment of the (complex) values of Γ_{12} and M_{12} . The relative argument between M_{12} and Γ_{12} is expected to be of $0(m_c^2/m_b^2)$ in the Standard Model. All together, a_{SL} is expected to be less than 10^{-3} for B_d^0 and less than 10^{-4} for B_S^0 and thus outside of the capabilities of the next future experimental facilities.

3.3 Rate Asymmetries

- By using the first method discussed in the last Section, one considers the process³⁵ $B \rightarrow J/\psi K_s$ able to measure $\sin(2\beta)$, where β is the angle of the unitarity triangle shown in Fig. 7. The quark diagrams responsible of this process are a tree diagram and a penguin diagram, as shown in Fig. 11




Figure 11: Tree and penguin diagrams relevant to the process $B \rightarrow J/\psi K_s$.

The penguin contribution is small, estimated to be less than 1% of the tree amplitude, and furthermore one discovers by inspection that both diagrams contain the same CP-phase, i.e., there is a unique (complex) hadronic amplitude to describe the exclusive process.

For B'_d s in the initial state, one has $\Delta\Gamma/\bar{\Gamma} \sim 5 \times 10^{-3}$, so $\Delta\Gamma$ can be neglected in the time distribution. The corresponding rate asymmetry for this definite final state $f \equiv J/\psi K_s$, eigenstate of CP, leads to

$$\frac{\Gamma_f - \bar{\Gamma}_f}{\Gamma_f + \bar{\Gamma}_f} = -\sin(2\beta) \sin(\Delta Mt)$$

independent of the (unique) hadronic amplitude !

This asymmetry is expected to be detected and measured at the Tevatron, the B -factories, HERA-B and at the LHC experiments. Precision determinations with $\sigma(\sin 2\beta) \sim 0.02$ can be expected.

Taking into account that the $(c\bar{c})$ system in the quark diagrams of Fig. 11 is, by itself, composed of charge-conjugated quarks, one can envisage the consideration of the semiinclusive process³⁶ $B \rightarrow K_s X(c\bar{c})$. No cancellation of the rate asymmetries is expected when summing over the different hadronic states of the $(c\bar{c})$ spectrum.

- With the aim to extract the angle α of the unitarity triangle, one considers the exclusive process $B_d \rightarrow \pi^+ \pi^-$, where $f \equiv \pi^+ \pi^-$ is again a CP-eigenstate.

The quark diagrams, shown in Fig. 12, contributing to the process are again a tree diagram and a penguin diagram




Figure 12: Tree and penguin diagrams relevant to the process $B_d \rightarrow \pi^+ \pi^-$.

The tree amplitude is governed by the quark mixing matrix element

$$V_{ub}^* V_{ud} \sim A\lambda^3(\rho + i\eta)$$

whereas the penguin amplitude contains

$$V_{tb}^* V_{td} \sim A\lambda^3(1 - \rho - i\eta)$$

so that both are of similar CKM strength. The effect of the loop and the gluonic exchange in the penguin amplitude has been estimated to give a relative contribution of the order of 10%-20%. An independent measure of these two contributions and their relative magnitudes is highly desirable.

Taking into account the contribution of the two different hadronic amplitudes, one induces a Time-Dependent CP-violating Rate Asymmetry

$$\frac{N - \bar{N}}{N + \bar{N}} = a \cos(\Delta mt) + b \sin(\Delta mt)$$

where "a" is obtained from the interference of the Direct CP violating Decay Amplitudes, whereas "b" comes from the interplay between Mixing and Decay.

Although the process $B_d \rightarrow \pi\pi$ is highly CKM-forbidden, the main problem is not statistics, particularly in the hadronic machines. Its major source of background comes from the accompanying decays $B_d \rightarrow K^+ \pi^-$; $B_s \rightarrow K^+ \pi^-, K^+ K^-$, which by themselves could have contributions to the asymmetry and not only to the rate. The virtues of having particle identification in the detector are evident, taking into account that: i) those backgrounds are reduced by Particle Identification, ii) the amount of background and its shape can be studied by reconstructing each decay mode using Particle Identification. These virtues are illustrated in Fig. 13, as advanced by the LHC-B collaboration




Figure 13: The virtues of Particle Id for $B_d \rightarrow \pi\pi$.

A purity of the data sample near 100% can be envisaged by reducing the efficiency to about 50%.

From the theoretical side, the absence of the penguin amplitude would lead to $a = 0$, $b = -\sin(2\alpha)$, where α as shown in the unitarity triangle of Fig. 7. The process $B_d \rightarrow \pi^+\pi^-$ is, however, theoretically difficult, because the presence of the penguin amplitude P affects the extraction of α as obtained from the tree amplitude T . To leading order in $|P/T|$ small, one has

$$\begin{cases} a = 2|\frac{P}{T}|\sin(\delta_P - \delta_T)\sin\alpha \\ b = -\sin(2\alpha) - 2|\frac{P}{T}|\cos(\delta_P - \delta_T)\sin\alpha\cos 2\alpha \end{cases}$$

where $\delta_{T(P)}$ is the elastic-final-state-interaction phase shift for the $T(P)$ hadronic amplitude.

If $|P/T|$ were known independently, the value of α could be extracted from the measurement of both "a" and "b". The quantity "b" is very sensitive to α and much less to the phase shift difference $\delta_P - \delta_T$, whereas the quantity "a" can determine $\delta_P - \delta_T$. The correlation between α and $\delta_P - \delta_T$ is quite small.

To determine $|P/T|$ experimentally, Gronau and London have used the isospin triangle relations³⁷ among the amplitudes M_{+0} for $B_u^\pm \rightarrow \pi^\pm\pi^0$, M_{+-} for $B_d \rightarrow \pi^+\pi^-$ and M_{00} for $B_d \rightarrow \pi^0\pi^0$

$$M_{+-} + \sqrt{2}M_{00} = \sqrt{2}M_{+0}$$

If all the three decay modes are measured, the value of $|P/T|$ can be extracted. However, $B_d \rightarrow \pi^+\pi^-$ is colour allowed in the quark diagram, whereas $B_d \rightarrow \pi^0\pi^0$ is colour suppressed and it needs a colour rearrangement. What is the price to be paid by this dynamics could be as high as to lower the branching ratio for $\pi^0\pi^0$ by a factor 10.

The channel $B_d \rightarrow \pi^+\pi^-$ could have other theoretical problems, like the existence of an electroweak (beyond the gluonic) penguin amplitude and/or inelastic-final-state-interaction effects. These problems need further scrutiny³⁸.

- With the strategy to show the consistency of the treatment, different methods to extract the angle $\gamma = \pi - \alpha - \beta$ from rate-asymmetries have been proposed. I mention here two of these proposals³⁹:

i) Use of four Time-Dependent Decay Rates from the B_s -meson

$$\begin{aligned} B_s \rightarrow D_s^- K^+ &\xleftrightarrow{CP} \bar{B}_s \rightarrow D_s^+ K^- \\ \bar{B}_s \rightarrow D_s^- K^+ &\longleftrightarrow B_s \rightarrow D_s^+ K^- , \end{aligned}$$

because the final state is not a CP-eigenstate in this case.

The Proper Time Distributions depend on the parameters x_s, Δ and γ , describing the mixing in the B_s -system, the final-state-strong-interaction phase and the relative weak CP-phase. The tree diagrams from B_s and $\bar{B}_s$ to a given final state are shown in the Fig. 14




Figure 14: Tree quark diagrams for B_s and $\bar{B}_s$ leading to $D_s^- K^+$.

Tree quark diagrams for B_s and $\bar{B}_s$ leading to $D_s^- K^+$

In order to resolve a twofold ambiguity in the extraction of γ and Δ from the experiment, an important role of a term like $\sinh(\Delta\Gamma_s t/2)$ is needed. For the B_s -system, $\Delta\Gamma_s/\bar{\Gamma}_s \sim 0.1$.

ii) Use of Decay to Self-Tagging final states, which are flavour-specific⁴⁰.

In this case one can use the rate asymmetries as for charged B's³⁴, with only rates and no time-dependence.

Take the six decays

$$\begin{aligned}
 B^0 &\longrightarrow D^0 \begin{cases} \longrightarrow K^+ \pi^- \\ \longrightarrow K^{*0} \\ \longrightarrow K^- \pi^+ \end{cases} + \text{c.c.} \\
 B^0 &\longrightarrow \bar{D}^0 \begin{cases} \longrightarrow K^+ \pi^- \\ \longrightarrow K^{*0} \\ \longrightarrow K^+ \pi^- \end{cases} + \text{c.c.} \\
 B^0 &\longrightarrow D_1 \begin{cases} \longrightarrow K^+ \pi^- \\ \longrightarrow K^{*0} \\ \longrightarrow K^+ K^-, \pi^+ \pi^- \end{cases} + \text{c.c.}
 \end{aligned}$$

where D_1 is the $CP = +$ eigenstate for $D^0 - \bar{D}^0$. If the corresponding amplitudes are indicated by $M, \bar{M}, M_+$, the ones for the conjugated processes are $\bar{M}, M, M_-$, respectively. The first three amplitudes close an isospin triangle

relation, and thus do the second three amplitudes. The relative orientation of the two triangles measures the angle (2γ), where γ is that shown in the unitarity triangle. This construction is illustrated in Fig. 15, where the amplitude M is taken as the reference in the complex plane.




Figure 15: The two isospin triangle relations for the six decays associated with $B \rightarrow DK^*$.

The two isospin triangle relations for the six decays associated with $B \rightarrow DK^*$

3.4 Outlook

The flavour structure is one of the main pending questions in fundamental physics. The Standard Model incorporates the Cabibbo-Kobayashi-Maskawa mechanism to generate CP-violation. For 3 families, one single CP-phase appears, leading to constrained physics. A fundamental explanation of the origin of CP-violation is however lacking.

The CP-odd asymmetries in the $B^0 - \bar{B}_0$ system can lead, by the choice of appropriate final states, to a determination of the unitarity triangle when combined with side measurements. The three angles α, β and $\gamma = \pi - \alpha - \beta$ can be separately determined and the experimental methods to do so have been discussed here. But still one perceives that all possible methods for the determination of the unitarity triangle have not yet been studied. The hadronic physics involved in some matrix elements needs further scrutiny. The overall consistency will need of different experimental methods to extract the sides and angles of the unitarity constraint.

References

1. G. Baur et al., PS 210 at CERN's LEAR, *Phys. Lett. B* **368**, 251 (1996).

2. T. D. Lee, C. N. Yang, *Phys. Rev.* **104** 254 (1956).
3. L. D. Landau, *Nucl. Phys.* **3** 127 (1957).
4. C. Y. Prescott et al., *Phys. Letters* **84B** 524 (1979).
5. M.A. Bouchiat, *Proc. Paris Neutral Currents* (1993), p. 247.
6. SLD Collaboration, K. Abe et al., *Phys. Rev. Lett.* **73**, 25 (1994).
7. The LEP Electroweak Working Group, CERN-PPE (1994).
8. J. Schwinger, *Phys. Rev.* **82** 914 (1951) ; G. Lüders, *Mat. Fys. Medd. Kongl. Dan. Vid. Selsk.* **28** (1954) No. 5. W. Pauli in *Niels Bohr and The Development of Physics*, Pergamon Press (1955)
9. J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, *Phys. Rev. Lett.* **13**, 138 (1964).
10. W. Grimus and M. N. Rebelo, UWTHPH-1995-7.
11. C. Jarlskog, in CP-Violation, *World Scientific* (1989), p. 3.
12. S. L. Glashow and S. Weinberg, *Phys. Rev. D* **15**, 1958 (1977). E.A. Paschos, *Phys. Rev. D* **15**, 1966 (1977).
13. S. L. Glashow, J. Iliopoulos and L. Maiani, *Phys. Rev. D* **2**, 1285 (1970).
14. J. Bernabéu and P. Pascual, *Electro-Weak Theory*, Univ. Auton. Barcelona, B-21399 (1981).
15. N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963).
16. M. Kobayashi and K. Maskawa, *Progr. Theor. Phys.* **49** 652 (1973).
17. Particle Data Book, Review of Particle Properties, *Phys. Rev. D* **54**, 1 (1996).
18. L. Wolfenstein, *Phys. Rev. Lett.* **51**, 1945 (1984).
19. M. Gell-Mann and A. Pais, *Phys. Rev.* **97**, 1387 (1955).
20. M. Bardon et al., *Ann. Phys. (NY)* **5**, 156 (1958); D. Neagu et al., *Phys. Rev. Lett.* **6**, 552 (1961).
21. J. M. Gaillard et al., *Phys. Rev. Lett.* **18**, 20 (1967); J. W. Cronin et al., *Phys. Rev. Lett.* **18**, 25 (1967).
22. S. Bennet et al., *Phys. Rev. Lett.* **19**, 993 (1967); D. Dorfman et al., *Phys. Rev. Lett.* **19**, 987 (1967).
23. L. Wolfenstein, *Phys. Rev. Lett.* **13**, 562 (1964)
24. P.P. Paulopoulos et al., CP-LEAR Collaboration, CERN-PPE-96-189, Workshop on K-physics, Orsay (1996).
25. G.D. Barr et al., NA31, *Phys. Lett. B* **317**, 233 (1993).
26. L.K. Gibbons et al., E731, *Phys. Rev. Lett.* **70**, 1203 (1993).
27. J. Bernabéu, F.J. Botella and J. Roldán, in CP-Violation in Particle Physics and Astrophysics, p. 389, ed. J. Tran Thanh Van, Editions Frontières (1990).
28. H. J. Lipkin, *Phys. Rev.* **176**, 1715 (1968).
29. I. Dunietz, J. Hauser and J.L. Rosner, *Phys. Rev. D* **35**, 2166 (1987).

- 30. J. Bernab  , F. J. Botella and J. Rold  n, *Phys. Lett. B* **211**, 226 (1988).
- 31. C. Jarlskog, *Phys. Rev. Lett.* **55**, 1039 (1985); *Z. Phys. C* **29**, 491 (1985); J. Bernab  , G. Branco and M. Gronau, *Phys. Lett. B* **169**, 243 (1986).
- 32. A. Ali and D. London, *Nuovo Cimento, Nuovo Cimento* **A109**, 957 (1996); A. Pich and J. Prades, *Phys. Lett. B* **346**, 342 (1995).
- 33. A. Carter and A.I. Sanda, *Phys. Rev. Lett.* **45**, 952 (1980); *Phys. Rev. D* **23**, 1567 (1981).
- 34. J. Bernab   and C. Jarlskog, *Z. Phys. C* **8**, 233 (1981).
- 35. I.I. Bigi and A. I. Sanda, *Nucl. Phys. B* **193**, 85 (1981).
- 36. J. Bernab   and C. Jarlskog, *Phys. Lett. B* **301**, 275 (1993).
- 37. M. Gronau and D. London, *Phys. Rev. Lett.* **65**, 3381 (1990).
- 38. G. Kramer and W. F. Palmer, *Phys. Rev. D* **52**, 6411 (1995).
- 39. R. Aleksan et al., ECFA 93-151 (1993).
- 40. M. Gronau and D. Wyler, *Phys. Lett. B* **265**, 172 (1991).